

# CHIROPRACTIC NEUROLOGY RESEARCH BRIEF

A QUARTERLY NEWSLETTER FOR HEALTH CARE PROFESSIONALS BY MARK SARACINO, DC, DACAN

VOLUME FOUR

ISSUE ONE

## Spinal Manipulative Therapy for Low Back Pain

### Introduction

Chiropractic manipulative therapy (CMT) has gained the attention of the medical and research communities, because substantial evidence has been found validating its efficacy for acute and the long-term treatment of low back pain (LBP). From 1993-2002, at least 43 randomized, controlled trials of spinal manipulation have been performed for acute, sub-acute and chronic LBP. 30 of these studies favor spinal manipulation over comparison treatments, 13 studies found no significant difference and not one study has found spinal manipulation to be statistically or clinically less effective than the comparison treatment. (1) CMT could be rising to be the most studied intervention for LBP.

Government-sponsored studies in the United States, Canada and New Zealand have recommended spinal manipulation for the treatment of LBP. (2-4) In 1994, the Agency for Health Care Policy and Research, a division of the U.S. Department of Health and Human Services, appointed a panel of experts to review over 3,000 studies on treatments for acute LBP and found that spinal manipulation was one of the recommended treatments. (2)

### Research Comparing CMT to Physical Therapy and Medicine

CMT has benefits beyond short-term relief of mild-to-moderate LBP. Comparing chiropractic and hospital outpatient treatment of LBP, the authors concluded that chiropractic treatment was more effective, especially for patients with chronic or severe pain. (5) At the three-year follow-up, the chiropractic group had a 29 percent advantage in Oswestry (disability) scores compared to the hospital group. (6)

A trial of 256 patients with non-specific low back and neck complaints compared spinal manipulation and physiotherapy (PT) of exercises, massage, heat, electrotherapy, ultrasound and shortwave diathermy, which I render, to that which a general practitioner rendered (analgesics, NSAIDs, advice about posture, home exercises, and bed rest) and placebo (detuned shortwave diathermy and ultrasound). CMT and PT were superior to the general practitioner and placebo groups, and at the 12-month follow-up, the manipulation group demonstrated greater improvement in the main complaint and physical functioning compared to PT. (7)

A 1998 *New England Journal of Medicine* study compared CMT, PT and an educational pamphlet for the treatment of LBP. Although the authors concluded that chiropractic and PT showed little benefit compared to the educational pamphlet, a closer look at the data found that the chiropractic group's subjects had worse histories, pain intensities and daily living and work activity risk factors. At the four and 12 week follow-ups, in spite of an unfavorable baseline, the chiropractic group fared better in terms of pain and disability and produced a significant decrease in the use of pain medication compared to the other groups (18 percent in the chiropractic group, 27 percent in the PT group, and 32 percent in the pamphlet group). At the 11-month follow-up, a smaller percentage of the chiropractic subjects lost time from work, required bed rest, and had limited daily living activity compared to the PT and pamphlet groups. (8)

A randomized controlled trial of one hundred fifteen patients with chronic spinal pain (greater than 13 weeks duration) were assigned to receive either NSAIDs (Celebrex, Vioxx, or paracetamol), acupuncture or CMT twice a week for nine weeks (9). By the end of the study, CMT achieved the best overall results with the highest proportion of patients receiving early and complete relief (27.3 percent) compared to acupuncture (9.4 percent) and medication (5 percent). The chiropractic group had a 50 percent improvement in back pain intensity compared to acupuncture (15 percent) and medication (0 percent) and neck and low back ranges-of-motion were dramatically increased. A notable finding is that the chiropractic patients had the highest average duration of chronic pain (8.3 years) versus the acupuncture group (6.4 years) and the medication group (4.5 years). One of the study's most remarkable finding was the chiropractic group reported a 47 percent improvement in "overall health" compared to 15 percent for the acupuncture group and 18 percent for the medication group.

### **CMT's Mechanism, Stabilization and Exercise Protocol**

CMT modulates pain at the zygapophyseal joints by causing a decrease in paraspinal hyperalgesia in subjects with symptoms and increase in paraspinal muscle pain thresholds in subjects with no symptoms. (10-13) CMT stimulates joint mechanoreceptors and inhibits second order neurons in the dorsal horn of the spinal cord which transmit nociception to the Limbic center of pain perception in the brain. (14)

CMT, also, stimulates muscle reflexes. (15-18) Colloca et al. found that 95 percent of subjects' mechanoreceptors of discoligamentous and muscular afferent nerve fibers were affected. (17) Herzog et al. states: "Because reflex pathways are evoked systematically during spinal manipulative treatment, there is a distinct possibility that these responses may cause some of the clinically observed beneficial effects, such as a reduction in pain and a decrease in hypertonicity of muscles." (16)

Another study investigated the effects of paraspinal muscle strength following spinal manipulative therapy. 40 subjects were assessed for trunk muscle strength by performing maximum voluntary contraction isometric trunk extensions while lying prone. The subjects received instrument-assisted SMT, sham manipulation or neither (control).

Spinal manipulation produced a significant increase in EMG erector spine isometric muscle output and trunk muscle strengths increased 21 percent. (19)

Several studies have shown that manipulation can increase spinal ranges-of-motion. (20-24) In Orthopedics, better lumbar sagittal (anterior to posterior) stability following lumbar spinal fusion produces better post-surgical outcomes. (25)

In Chiropractic, lumbar extension traction has been shown to increase the lumbar lordosis (normal, forward-arching curve) in chronic LBP patients with hypolordosis (flattening of the forward curve). (26) 48 patients with chronic LBP were matched for sex, age, height, weight, and pain scores with 30 control subjects. The treatment group received lumbar spinal manipulation in the first few weeks for short-term pain relief, then sessions of 3-point lumbar extension traction three to four times per week for 12 weeks. Pain scores and radiographic measurements did not change in the control group, but in the extension traction group, pain scores decreased from 4.4 to 0.6 on a scale of 1-10.

Radiography showed statistical improvements in lumbar lordotic and sacral angles increases. 34 of 38 of the subjects were evaluated for 17.5 months long-term follow-up and ALL 34 subjects maintained an improved lumbar lordosis. This chiropractic method of lumbar lordosis exercising appears to be the first conservative procedure to increase the lordosis in chronic LBP patients for an extended period of time.

## **Discussion**

Improving “overall health” from better spinal functioning by balancing loads which decreases aberrancy on spinal tissues is a paradigm of Chiropractic Neurology’s practice guidelines and philosophy. CMT has been found to be an effective treatment for low back pain in numerous studies and should be recommended for patients with both acute and chronic low back pain.

## **REFERENCES**

1. Meeker WC, Haldeman S. Chiropractic: a profession at the crossroads of mainstream and alternative medicine. *Annals Internal Medicine* 2002; 136:216-27.
2. Bigos S, et al. Acute low back problems in adults. Clinical practice guideline no. 14. AHCPR Pub No. 95 – 0624. Rockville, MD: Agency for Health Care Policy and Research, Public Health Service, U.S. Dept. of Health And Human Services. December 1994.
3. Manga P. The effectiveness and cost effectiveness of chiropractic management of low back pain. Ottawa, Ontario, Canada. Ontario Ministry of Health 1993.
4. Chiropractic in New Zealand. Report of the Commission of Inquiry. Wellington, New Zealand: PD Hasselberg, Government Printer, 1979.
5. Meade TW et al. Low back pain of mechanical origin: randomized comparison of chiropractic and hospital outpatient treatment. *British Medical Journal* 1990; 300: 1431-37.
6. Meade TW et al. Randomized comparison of chiropractic and hospital outpatient management for low back pain: results from extended follow-up. *British Medical Journal* 1995; 311: 349-51.
7. Koes BW et al. Randomized clinical trial of manipulative therapy and physiotherapy for persistent back and neck complaints: results of one year follow-up. *British Medical Journal* 1992; 304:601-605.
8. Cherkin DC et al. A comparison of physical therapy, chiropractic manipulation and provision of an educational booklet for the treatment of patients with low back pain. *New England Journal Medicine* 1998; 339: 1021-29.
9. Giles LG, Muller R. Chronic spinal pain: A randomized clinical trial comparing medication, acupuncture, and spinal manipulation. *Spine* 2003; 28:1490-1502.
10. Hsieh J, Hong CZ. The effect of chiropractic manipulation on the pain threshold of myofascial trigger points: a pilot study. In: Wolk S, editor. *Proceedings of the 1990 International Conference on Spinal Manipulation*. Foundation for Chiropractic Education and Research, Washington D.C.; 1990.

11. Vernon HT et al. Pressure pain threshold evaluation of the effect of spinal manipulation in the treatment of chronic neck pain: a pilot study. *Journal Manipulative Physiological Therapeutics* 1990; 13:13-6.
12. Vernon HT et al. Evaluation of neck muscles strength with a modified sphygmomanometer dynamometer: reliability and validity. *Journal Manipulative Physiological Therapeutics* 1992; 15:34-9.
13. Terret ACJ, Vernon HT. Manipulation and pain in tolerance: a controlled study of the effect of spinal Manipulation on paraspinal cutaneous pain tolerance levels. *American Journal Physical Medicine* 1984; 63: 217-25.
14. Vernon HT. Qualitative review of studies of manipulation-induced hypoalgesia. *Journal Manipulative Physiological Therapeutics* 2000; 23: 134-8.
15. Pickar JG, Wheeler JD. Response of muscle proprioceptors to spinal manipulative-like loads. *Journal Manipulative Physiological Therapeutics* 2001; 24: 2-11.
16. Herzog et al. Electromyographic responses of back and limb muscles associated with spinal manipulative therapy. *Spine* 1999; 24 (2): 146-52.
17. Colloca CJ, Keller TS. Electromyographic reflex responses to mechanical force, manually assisted spinal adjustments. *Spine* 2001; 26(10): 1117-24.
18. Colloca CJ, Keller TS. Stiffness and neuromuscular reflex response of the human spine to dynamic posteroanterior manipulative thrusts in patients with low back pain. *Journal Manipulative Physiological Therapeutics* 2001; 24: 489-500.
19. Colloca CJ, Keller TS. Mechanical force spinal adjustment increases trunk muscle strength assessed by EMG: A comparative controlled clinical trial. *Journal Manipulative Physiological Therapeutics* 2000; 23: 585-95.
20. Cassidy JD et al. The effect of manipulation on pain and range of motion in the cervical spine. *Journal Manipulative Physiological Therapeutics* 1992; 15: 495-500.
21. Cassidy JD et al. The immediate effect of manipulation vs. mobilization on pain and range of motion in the cervical spine: a randomized controlled trial. *Journal Manipulative Physiological Therapeutics* 1992; 5: 570-5.
22. Hviid H. The influence of chiropractic treatment on the rotary mobility of the cervical spine. *Ann Swiss Chiropractic Assoc* 1971; 5:31-44.
23. Nilsson N et al. Lasting changes in passive range of motion after spinal Manipulation: a randomized, blind, controlled trial. *Journal Manipulative Physiological Therapeutics* 1996; 19:165-8.
24. Lehman GJ, McGill SM. Spinal manipulation causes variable spine kinematic and his trunk muscle electromyographic responses. *Clinical Biomechanics (Bristol, Avon)* 2001; 16 (4): 293-9.
25. Kawakami M et al. Lumbar sagittal balance influences the clinical outcome after decompression and posterolateral spinal fusion for degenerative lumbar spondylolisthesis. *Spine* 2002; 27 (1): 59-64.
26. Harrison DE et al. Changes in sagittal lumbar configuration with a new method of extension traction: nonrandomized clinical controlled trial. *Archives Physical Medicine Rehabilitation* 2002; 83 (11): 1585-91.

*Mark Saracino, Board Certified Chiropractic Neurologist  
Diplomat American Chiropractic Academy of Neurology  
(adjacent to the Valley Forge Convention Center and Radisson Hotel)  
1150 First Avenue, suite 120  
King of Prussia, Pennsylvania 19406 1341*

*voice 610 337 3335  
fax 610 337 4858  
www.ACNB.org  
Mark.Saracino1@juno.com  
www.DrSaracino.com*

*(see back issues at "Health Information" link)*